

Onondaga County Health Department

**Division of Environmental Health
421 Montgomery Street
Syracuse, New York 13202**

Incinerator Monitoring Program

2012 Screening Summary for Organic Constituents

June 1, 2013

Submitted To: Cynthia B. Morrow, M.D., M.P.H.
Commissioner of Health

Submitted By: Kevin L. Zimmerman
Director, Division of Environmental Health

Contents:

- I. Table of Abbreviations.
- II. Executive Summary.
- III. Introduction.
- IV. PCDD/PCDF Specific Discussion.
- V. PCB Specific Discussion.
- VI. PAH Specific Discussion.
- VII. Summary and Conclusions.
- VIII. Routine Soil Sample Site Location Map.

List of Tables

- Table 1: PCDD/PCDF Results; Control and Routine Site Soils
- Table 2: PCDD/PCDF Results; WTE Facility Combined Ash
- Table 3: PCB Results; Control and Routine Site Soils
- Table 4: PCB Results; WTE Facility Combined Ash
- Table 5: PAH Results; Control and Routine Site Soils
- Table 6: PAH Results; WTE Facility Combined Ash

Attachment A: Historical PCDD/PCDF Results

Attachment B: Historical PCB Results

Attachment C: PAH Background Soil Concentrations

I. Table of Abbreviations

The following abbreviations may be used in this report:

ATSDR	Agency for Toxic Substances and Disease Registry.
PCDD/PCDF	Polychlorinated Dibenzo-p-Dioxins/Dibenzofurans.
PCB	Polychlorinated Biphenyls.
PAH	Polycyclic Aromatic Hydrocarbons.
µg/g	micrograms per gram (also denoted as ug/g).
ng/g	nanograms per gram.
ng/kg	nanograms per kilogram (pg/g equivalent).
pg/g	picograms per gram (ng/kg equivalent).
LD	Limit of Detection.
NS	Not sampled.
ND	Not detected.
OCHD	Onondaga County Health Department.
WTE	Waste to Energy.
~	approximately.
<	Less than.
>	Greater than.

II. Executive Summary

Organic sample analyses for the year 2012 of soil and ash for the Incinerator Monitoring Program have been conducted by Axys Analytical Services LTD. Analyses for this summary include PCDD/PCDF, PCB and PAH. Ash collection was conducted by Covanta Energy personnel (formerly Odgen Martin), with random oversight by the Onondaga County Health Department's Division of Environmental Health. The collection of all ambient environmental samples was, and continues to be, the responsibility of the Division of Environmental Health. Final sample composites were prepared by Life Science Laboratories, Inc. (formerly O'Brien and Gere Environmental Laboratory).

Much of the comparative background data and information referenced in this report was obtained from NYDEC Soil Cleanup Objectives, EPA Preliminary Remediation Goals and Soil Screening Levels, along with the U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry's Toxicological Profiles.

The analyses for organics completed during this monitoring period show the parameters are within the expected range for urban and rural environments. The results are below levels associated with health risk. The 2012 sampling revealed levels typical of historical data at all sites. Given the low levels detected and the corresponding variation expected as a result of sample collection, preparation, and laboratory procedures, the levels that have been determined do not allow for comparison to establish change in the environment. In the organic monitoring conducted to date, no relationship between the operation of the incinerator and increased levels of organics has been established.

III. Introduction.

In November 1994, the Onondaga County Resource Recovery Agency, in contract with the Covanta Energy Company (formerly Ogden Martin Company), commenced operation of a municipal solid waste incinerator. This undertaking was part of a multifaceted solid waste management program to achieve a reduction of volume of landfill waste, energy withdrawal and the removal of solids incompatible with incineration. Part of the management program for the reuse of materials and the removal of materials prior to the municipal waste stream had been started earlier.

The Onondaga County Health Department initiated a program in 1993 to include short and long term monitoring aspects to document any health implications to the public and environmental changes from the incinerator. Changes have been made to the monitoring program several times over the course of time in response to new information as it became available. In 2003 the monitoring program was re-evaluated to provide a more effective and efficient program. Direct interaction was established with the Onondaga County Resource Recovery Agency (OCCRA) and the New York State Department of Environmental Conservation (DEC) in providing stack monitoring results and improved assurance on reporting of adverse events and equipment failures. This allowed for effective evaluation of short-term change in the incinerator emissions rather than the previous limited scope offsite air monitoring conducted over a nine year period. Several changes were implemented in 2009 based on the low levels of organic constituents detected in the monitoring conducted to date, and the fact that there is no evidence of a trend or levels associated with health risks. The fourteen routine soil sites (which include two control sites) continue to be sampled and analyzed twice a year for metals which are documented in a separate report. Half of the sites (7, including one control) are being tested for organics once a year. The sites will be rotated so that each is tested every other year. The program includes the flexibility to test a site two years in a row if there is an elevated level of any organic constituent. The four ash route soil sites have been eliminated from the program. These sites were located along the route that trucks take to carry ash across and out of the County. To date these sites have not shown any elevation of metals or organics and the trucks are covered at all times. Ash, directly from the incinerator continues to be analyzed for metals twice a year and organics once a year. The department continues to interact directly with OCCRA and DEC in review of stack monitoring results.

This is the thirteenth report for screening of organics, analyzed for dioxin, dibenzofurans, polychlorinated biphenyls and polycyclic aromatic hydrocarbons, from samples of ambient soil and combined ash collected from the incinerator operation. The analysis of soil samples provides a useful and convenient mechanism for monitoring accumulative change of these organics in the environment. Surface soil samples can be utilized to monitor deposition of transient materials that can drop from atmospheric particulate materials, materials spilled in the area and materials spread on the land for agricultural purposes.

A program designed to monitor soil samples collected on a routine basis will provide an assessment of the organic material deposited in the sample area. The limitation of this matrix is that there are numerous sources and a normal action by nature is occurring on the soil at all times. The results reported should be utilized with other reports for studies in other areas. The soil sample analyses described in this

report are part of an ongoing program of environmental monitoring performed by the Onondaga County Health Department as part of its overall Incinerator Monitoring Program.

This report represents data from the screening of soil and ash collected during the calendar year 2012. This is the eighteenth year of operation of the WTE facility. Three samples were collected at each soil location during each sampling event. Ash sampling is conducted by Covanta Energy personnel during their semi-annual collection. Through the sampling year 2002, it was the responsibility of the Onondaga County Health Department Environmental Toxicology Laboratory to create the soil and ash composites. Beginning with 2004, the contract laboratory, Life Science Laboratories, Inc. (formerly O'Brien and Gere Environmental Laboratories) created one composite sample for each organic analysis of soil and a two-day and three-day composite of the ash for analysis.

IV. PCDD/PCDF Specific Summary.

PCDDs are a class of chlorinated tricyclic aromatic hydrocarbons. There are 75 chlorinated dioxins, all varying in toxicity. Generally, the PCDD congeners of relative toxic concern are 2,3,7,8-TCDD, 1,2,3,7,8-PeCDD, 1,2,3,4,7,8-HxCDD, 1,2,3,6,7,8-HxCDD, 1,2,3,7,8,9-HxCDD and 1,2,3,4,6,7,8-HpCDD. PCDFs are also a class of chlorinated tricyclic aromatic hydrocarbons. There are 135 chlorinated furans, of which, approximately 10 to 12 are expected to have significant acute toxicity. The most acutely toxic isomers appear to be 2,3,7,8-TCDF, 1,2,3,7,8-PeCDF and 2,3,4,7,8-PeCDF. Each sample was tested for seventeen different congeners of PCDD/PCDF.

Each congener of PCDD/DF has associated with it a toxic equivalency factor, TEF. This factor is an indication of the toxicity of the individual congeners with respect to 2,3,7,8 TCDD, the most toxic congener. Each sample has a calculated total toxic equivalency, TEQ, shown in Tables 1 and 2. While the toxic equivalency is the main consideration for the determination of change, individual congener concentrations have also been reviewed for significance.

Table 1 displays the results of soil analyses for dioxin and dibenzofurans at the six routine sampling sites and one control site for the spring sampling period of year 2012. In general, the set of TEQ results from these samples confirmed very well the results that were presented in the previously issued "Screening Summary for Organic Constituents" reports (Refer to Attachment A). Results from both the routine sites and the control site demonstrate no distinct pattern from background through year 2012 sampling. The TEQ's for 2012 are well below the screening level of 50 pg/g used by ATSDR and the EPA preliminary remediation goal of 1000 pg/g. The levels as reported are not of health significance and are within expected levels as stated in other documents for background levels in soil.

Table 2 displays the analyses for ash from the incinerator. The TEQ result for the ash composite for day 1 & 2 and day 3-5 are consistent with previous results. Ash is not homogenous and can contain chunks of material which may account for an occasionally inconsistent result. These results are similar to reports for ash identified by other investigators and reported in published literature. All of the ash is transported in closed vehicles and buried at a Department of Environmental Conservation permitted

landfill.

Attachment A shows the historical TEQ values for routine soil sites, control sites, and ash samples.

V. PCB Specific Summary.

Polychlorinated biphenyls, PCB's, are a class of more than 200 man-made chemical compounds. PCB's were widely used in industrial applications due to the physical characteristics of the compounds. Incineration of PCB containing products can lead to a release of PCB's into the environment. Soil sampling is a strong indicator of PCB levels in the environment because of the persistence and adhesion capabilities of the substance. PCB analysis in the past had resulted in less than detectable concentrations. Axys Analytical Services, LTD lowered the limit of detection for PCB starting in 2000 so that usable concentrations are now being presented. The ATSDR Toxicological Profile for PCB (1996) indicates that typical mean PCB concentrations in background soil are <100 µg/kg (<100,000 pg/g). The NYSDEC has a Soil Cleanup Objective of 100,000 pg/g for PCB's and the EPA has a soil screening level of 240,000 pg/g for residential soil.

PCB results are presented in Tables 3 and 4. Table 3 displays results for the six routine sites within the impact area of the WTE Facility along with a control site. The mean PCB concentration for routine sites was 4,255 pg/g, with a maximum concentration of 17,400 pg/g at the Syracuse University site. In general, PCB results in this study are well below the ATSDR typical background soil level of 100,000 pg/g. Soil is not homogenous and can contain materials that can account for an occasional inconsistent result. Attachment B shows historical levels of PCB's at routine soil sites along with control sites.

Table 4 displays the results of PCB analyses of ash as collected from the WTE Facility. At 1,800 and 20,500 pg/g, the PCB levels for the year 2012 sampling are lower than the previously stated ATSDR typical background soil level.

VI. PAH Specific Summary.

Polycyclic aromatic hydrocarbons, PAH's, are primarily formed as the result of incomplete combustion of organic matter. PAH's, like PCB's, have a strong persistence and affinity to particulate matter. For this reason, soil and ash sampling are quality measures of the levels attributable to incineration. As with the PCB analyses, Axys Analytical Services, LTD has lowered the limit of detection for PAH congeners for this report so that additional usable concentrations are now being presented.

PAH results for soil are presented in Table 5. Attachment C presents NYSDEC Soil Cleanup Objectives, EPA screening levels, NYS Rural soil survey results, and Toxicological Profile levels for PAH's for rural, agricultural and urban soils. These levels can vary widely for the individual PAH's. The levels reported in the 2012 study are generally within these expected ambient levels.

PAH results for the WTE ash composites are presented in Table 6. Comparison

of the 2012 composite ash results to the averages for years 1999 through 2011 individual results exhibits little variation in PAH congener specific concentrations.

VII. Summary and Conclusions

This screening represents the organic analysis data for calendar year 2012 environmental soil and ash samples. PCDD, PCDF, PCB and PAH levels are all quality indicators of ambient conditions in the environment. By following the concentrations and trends of these compounds, two objectives are accomplished. First, ambient conditions are monitored for changes due to point sources. Second, health risks can be established for the effect of the soil concentrations.

The reported concentrations of all organic compounds in this screening are within expected levels and are below significant health risk levels. In general, little change in levels of these compounds have been observed from background through the present organic screening period.

The Onondaga County Health Department will continue to monitor soil and ash for organic compounds.

OCHD ROUTINE SOIL MONITORING SITES




Table 1

		Control Site		Routine Sites					
		Dutch Hill	Clark Reservation	Jamesville Beach	Nob Hill	Sentinel Heights	Syracuse University	J.D. High School	
		15-May-12	15-May-12	15-May-12	15-May-12	15-May-12	15-May-12	15-May-12	
Accession Number:		L18216-5	L18216-7	L18216-6	L18216-1	L18216-2	L18216-4	L18216-3	
PCDD / PCDF	TEF								
2,3,7,8-TCDD	1	K 0.135	0.205	K 0.075	0.095	0.612	K 0.148	K 0.176	
1,2,3,7,8-PeCDD	0.5	0.456	0.496	0.184	0.252	4.31	0.565	0.306	
1,2,3,4,7,8-HxCDD	0.1	K 0.650	0.559	0.216	0.288	7.85	0.797	0.308	
1,2,3,6,7,8-HxCDD	0.1	1.16	1	0.445	0.565	16.3	1.41	0.62	
1,2,3,7,8,9-HxCDD	0.1	1.82	1.26	0.513	0.676	21.4	2.21	0.787	
1,2,3,4,6,7,8-HpCDD	0.01	13.9	19.5	7.94	10	385	23.5	11.7	
OCDD	0.001	64.1	114	43.1	67.4	2500	150	66	
2,3,7,8-TCDF	0.1	1.76	1.47	0.47	0.679	0.667	0.989	0.776	
1,2,3,7,8-PeCDF	0.05	0.653	0.515	K 0.217	0.243	0.722	0.281	0.339	
2,3,4,7,8-PeCDF	0.5	1.56	0.613	K 0.246	0.343	0.68	0.411	0.408	
1,2,3,4,7,8-HxCDF	0.1	2.19	0.937	0.398	0.488	4.15	0.769	0.682	
1,2,3,6,7,8-HxCDF	0.1	1.93	0.733	0.233	0.336	2.95	0.676	0.467	
1,2,3,7,8,9-HxCDF	0.1	0.107	K 0.085	< 0.0493	< 0.0488	0.189	< 0.0483	< 0.0498	
2,3,4,6,7,8-HxCDF	0.1	3.15	0.709	0.293	0.381	2.12	0.475	0.442	
1,2,3,4,6,7,8-HpCDF	0.01	11.2	5.97	3.41	3.03	73.4	5.33	4.07	
1,2,3,4,7,8,9-HpCDF	0.01	0.649	0.48	0.288	0.333	4.6	0.381	0.348	
OCDF	0.001	5.43	8.14	6.22	4.53	166	7.8	5.21	
Total TEQ		2.47	1.75	0.488	0.488	16	1.63	0.951	
Total Tetra-Dioxins		1.41	3.42	0.984	1.37	4.15	1.3	1.78	
Total Penta-Dioxins		5.84	4.48	1.39	1.24	18.2	2.95	2.72	
Total Hexa-Dioxins		13	10.7	4.62	5.72	157	15	5.77	
Total Hepta-Dioxins		27.9	37.8	14.4	19.1	808	48.5	22.6	
Total Tetra-Furans		8.66	11.3	2.49	3.72	5.82	7.06	5.3	
Total Penta-Furans		16.2	8.32	1.89	3.26	14.6	5.69	4.97	
Total Hexa-Furans		23	9.24	3.82	4.88	76.3	7.94	6.08	
Total Hepta-Furans		17	11.3	7.44	5.78	226	10.4	7.25	

Results reported in pg/g (ng/kg equivalent) dry weight.

Comparison Values

EPA Action Level	1,000 ng/kg (Total TEQ)
ATSDR EMEG Value	50 ng/kg (Total TEQ)

Table 2

		Combined Ash	
		Spring 2012	
		Day 1 & 2	Day 3 - 5
Accession Number:		L18216-8	L18216-9
PCDD / PCDF	TEF		
2,3,7,8-TCDD	1	14.2	14.7
1,2,3,7,8-PeCDD	0.5	30.6	33.3
1,2,3,4,7,8-HxCDD	0.1	16.9	21.4
1,2,3,6,7,8-HxCDD	0.1	25.3	35.3
1,2,3,7,8,9-HxCDD	0.1	39.6	53.1
1,2,3,4,6,7,8-HpCDD	0.01	184	276
OCDD	0.001	527	786
2,3,7,8-TCDF	0.1	210	224
1,2,3,7,8-PeCDF	0.05	94.2	96.7
2,3,4,7,8-PeCDF	0.5	76.4	83.2
1,2,3,4,7,8-HxCDF	0.1	88.3	80.9
1,2,3,6,7,8-HxCDF	0.1	91	94.7
1,2,3,7,8,9-HxCDF	0.1	7.62	8.25
2,3,4,6,7,8-HxCDF	0.1	50.5	62.4
1,2,3,4,6,7,8-HpCDF	0.01	203	238
1,2,3,4,7,8,9-HpCDF	0.01	26.1	29.7
OCDF	0.001	109	108
Total TEQ		116	127
Total Tetra-Dioxins		216	221
Total Penta-Dioxins		277	300
Total Hexa-Dioxins		312	401
Total Hepta-Dioxins		386	576
Total Tetra-Furans		1780	1850
Total Penta-Furans		1280	1340
Total Hexa-Furans		749	805
Total Hepta-Furans		312	364

Results reported in pg/g dry weight.

Table 3

	Control Site		Routine Sites										
	Dutch Hill 15-May-12		Clark Reservation 15-May-12		Jamesville Beach 15-May-12		Nob Hill 15-May-12		Sentinel Heights 15-May-12		Syracuse University 15-May-12		J-D High School 15-May-12
Accession Number:	L18216-5		L18216-7		L18216-6		L18216-2		L18216-3		L18216-5		L18216-3
PCB													
Total Monochloro Biphenyls	1.09		7.43		1.53		1.92		1.74		9.38		2.48
Total Dichloro Biphenyls	4		14.8		3.99		9.99		4.48		59.5		9.51
Total Trichloro Biphenyls	15.3		42.4		9.97		18.9		14.2		208		26.8
Total Tetrachloro Biphenyls	106		146		29.5		66.2		63.2		490		87.8
Total Pentachloro Biphenyls	207		467		122		300		254		1610		261
Total Hexachloro Biphenyls	741		850		246		657		418		6410		452
Total Heptachloro Biphenyls	811		785		163		461		307		6470		373
Total Octachloro Biphenyls	368		402		94.8		216		148		1920		178
Total Nonachloro Biphenyls	72.3		127		27.2		64.4		36.2		177		49
Decachloro Biphenyl	26.6		43.2		8.95		26.4		14.1		21.5		12.9
Total PCB'S	2350		2890		707		1820		1260		17400		1450

Results reported in pg/g dry weight.

Comparison Value	
ATSDR Typical Mean Background Value	< 100,000 pg/g

Table 4

	Combined Ash	
	Spring 2012	
	Day 1 & 2	Day 3 - 5
Accession Number:	L18216-8	L18216-9
PCB		
Total Monochloro Biphenyls	122	183
Total Dichloro Biphenyls	143	4260
Total Trichloro Biphenyls	240	6920
Total Tetrachloro Biphenyls	334	5660
Total Pentachloro Biphenyls	419	1600
Total Hexachloro Biphenyls	255	865
Total Heptachloro Biphenyls	161	662
Total Octachloro Biphenyls	72.7	298
Total Nonachloro Biphenyls	29.7	53.8
Decachloro Biphenyl	20.2	27.1
Total PCB'S	1800	20500

Results reported in pg/g dry weight.

Table 5

	Control Site		Routine Sites										
	Dutch Hill 15-May-13		Clark Reservation 15-May-12		Jamesville Beach 15-May-12		Nob Hill 15-May-12		Sentinel Heights 15-May-12		Syracuse University 15-May-12		J.D. High School 15-May-12
Accession Number:	L18216-5		L18216-7		L18216-6		L18216-1		L18216-2		L18216-4		L18216-3
PAH													
Naphthalene	2.52		14.4		3.35		4.17		13.7		4.24		3.99
Acenaphthylene	2.13		47.5		3.02		9.42		43.6		5.54		6.6
Acenaphthene	0.888		9.4		0.938		3.16		3.48		1.75		1.97
Fluorene	0.509		8.47		0.45		2.63		1.8		2.41		0.9
Phenanthrene	11.6		217		8.82		64.1		42.9		33		25
Anthracene	2		40		1.86		11.5		27.3		6.48		6.4
Fluoranthene	20.2		379		15.7		156		119		68.7		60.7
Pyrene	17.8		331		14.1		135		129		58.6		53.7
Benzo(A)Anthracene	7.1		134		6.24		50.2		73.9		24		27.6
Chrysene	12.5		212		10.9		76.3		110		41.5		40.3
Benzo(B,J,K)Fluoranthene	20		339		18.9		119		213		62.8		70.2
Benzo(E)Pyrene	8.33		132		7.78		48		91.7		25.2		26.3
Benzo(A)Pyrene	14.6		176		10.8		65.9		133		31.2		39.6
Perylene	1.88		32.4		1.56		14.4		27.7		6.71		7.91
Dibenzo(A,H)Anthracene	K 1.73		27.8		K 1.64		10.2		20.3		5.41		6.54
Indeno(1,2,3-CD)Pyrene	9.24		132		7.98		48.4		95.9		23.9		27.8
Benzo(G,H,I)Perylene	8.4		121		7.93		45.1		94.6		23.4		27.8
2-Methylnaphthalene	2.45		14.8		3.51		3.91		7.12		5.25		3.38
2-Chloronaphthalene							0.06				0.062		

Results reported in ng/g dry weight.

Table 6

	Combined Ash	
	Spring 2012	
	Day 1 & 2	Day 3 - 5
Accession Number:	L18216-8	L18612-9
PAH		
Naphthalene	71.3	107
Acenaphthylene	24.7	29.7
Acenaphthene	220	328
Fluorene	101	63.1
Phenanthrene	131	382
Anthracene	27.6	71.9
Fluoranthene	135	408
Pyrene	107	374
Benzo(A)Anthracene	51.9	190
Chrysene	60.9	215
Benzo(B,J,K)Fluoranthene	90.1	335
Benzo(E)Pyrene	272	240
Benzo(A)Pyrene	61.2	227
Perylene	14.1	64.7
Dibenzo(A,H)Anthracene	7.55	34.5
Indeno(1,2,3-CD)Pyrene	34.6	160
Benzo(G,H,I)Perylene	33.8	163
2-Methylnaphthalene	23.2	37.7
2-Chloronaphthalene	0.857	0.797

Results reported in ng/g dry weight.

Dioxin/Furan TEQ Soil Results through Year 2012 (pg/g dry weight)

Routine Soil Sites

Site	Year													
	1994	1999	2000	2001	2002	2004	2005	2006	2007	2008	2009	2010	2011	2012
Clark Reservation	1.8	1.2	2.27	1.42	1.23	2.03	1.90	1.76	1.73	1.26	***	1.64	***	1.75
Jamesville Beach	0.6	0.5	1.09	0.82	0.70	0.71	0.97	0.86	0.93	0.77	***	0.52	***	0.488
OCCF	0.79	2.2	1.68	1.47	1.26	1.38	5.54	1.52	1.94	1331.72@	1.72	***	2.13	***
DOT @ Jaquith	2		1.5	1.64	3.41	2.41	3.78	3.38	1.73	39.90@	2.62	***	3.95	***
Dutch Hill *	0.77		1.41	1.16	1.40	1.03	1.26	1.02	1.02	0.64	***	0.73	***	2.44
Erie - Poolsbrook*	1.39		1.5	1.14	1.86	**	**	**	**	**	**	**	**	**
Nottingham	0.51		0.78	0.79	0.80	0.70	0.94	0.85	0.84	0.74	0.76	***	0.43	***
SHFD	12		8.02	9.89	9.72	7.02	8.09	6.27	7.20	10.74	***	7.12	***	16
Sevier Rd	1.8		2.07	2.58	2.56	**	**	**	**	**	**	**	**	**
Beaver Lake *			0.51	0.53	0.85	0.70	0.72	0.64	0.69	0.65	0.38	***	0.5	***
Ch. 3 Towers			3.36	3.88	3.35	9.66	7.79	7.69	5.39	2.44	3.72	***	0.45	***
Gen.Crushed Stone			2.77	1.98	2.13	**	**	**	**	**	**	**	**	**
Highland Forest			1.18	1.24	0.96	**	**	**	**	**	**	**	**	**
JD High School			1.32	1.29	1.12	1.10	1.48	1.16	1.06	1.28	***	1.13	***	0.951
Nob Hill			0.93	0.91	0.90	6.83	1.01	1.00	1.07	1.05	***	0.78	***	0.488
Pratts Falls			0.91	0.98	0.77	0.87	0.98	0.83	0.94	1.17	0.82	***	0.94	***
Southwood			0.6	1.14	1.01	1.08	1.05	0.97	1.09	1.01	0.80	***	0.93	***
Syracuse University			3.11	6.97	9.47	13.89	3.14	3.66	12.96	0.67	***	2.45	***	1.63

* Denotes Control Sites

** Site no longer sampled due to program re-evaluation

*** Site not sampled this year. Sites are sampled every other year.

@ A single elevated value will not be assumed to be indicative of a change at a specific site, rather a pattern of values must demonstrate a statistically significant difference.

Combined Ash

Site	Year													
	1999-Spring	1999-Fall	2000-Fall	2001-Fall	2002-Fall	2004-Spring	2005-Spring	2006-Spring	2007-Spring	2008-Spring	2009-Spring	2010-Spring	2011-Spring	2012-Spring
Day 1 and 2	256	153	109	123	177	72	191	246	250	243	168	200	197	116
Day 3, 4, and 5	242	205	154	137	220	445	142	148	276	240	126	172	129	127

Note: For reference purposes, the ATSDR investigation level for Dioxin/Furan TEQ is 50 pg/g and the EPA clean up level is 1,000 pg/g.

Attachment B

PCB Results through Year 2012 (pg/g dry weight)

Routine Soil Sites

Site	2000	2001	2002	2004	2005	2006	2007	2008	2009	2010	2011	2012
Clark Reservation	6010	2360	3150	2780	3610	2770	4110	2640	***	2960	***	2980
Jamesville Beach	1260	644	683	703	1110	781	1220	1610	***	589	***	707
OCCF	3080	5230	2000	2310	6940	3120	6320	2190	2810	***	2650	***
DOT @ Jaquith	16100	15400	45100	9220	67100	49100	18000	14200	34700	***	31800	***
Dutch Hill *	2210	1170	1400	1200	1380	1140	1450	1340	***	1060	***	2350
Erie - Poolsbrook *	2620	1400	2020	**	**	**	**	**	**	**	**	**
Nottingham	2140	2280	3610	1640	7380	2850	3050	2110	4200	***	2020	***
SHFD	3080	2970	1760	1900	2730	1610	2510	1730	***	2240	***	1260
Sevier Rd	1870	1600	2250	**	**	**	**	**	**	**	**	**
Beaver Lake *	1970	1210	5250	2650	1420	1360	1360	1370	2450	***	1110	***
Ch. 3 Towers	3360	2310	2490	1620	1830	1730	2220	1400	1510	***	723	***
General Crushed Stone	9430	3160	5450	**	**	**	**	**	**	**	**	**
Highland Forest	2120	1210	1270	**	**	**	**	**	**	**	**	**
JD High School	3580	1780	1732	1810	2640	1780	1720	2720	***	1750	***	1450
Nob Hill	3500	2480	2500	3440	2810	2970	2830	2950	***	2510	***	1820
Pratts Falls	1890	1840	1440	1620	1650	1220	1450	2050	1230	***	1910	***
Southwood	2240	2160	1150	1480	1470	1470	2750	1640	1640	***	1120	***
Syracuse University	10700	114000	11000	9510	6940	11400	10900	1170	***	78600	***	17400

* Denotes Control Sites

** Site no longer sampled due to program re-evaluation

*** Site not sampled this year. Sites are sampled every other year.

Combined Ash

Site	2000-Fall	2001-Fall	2002-Fall	2004-Spring	2005-Spring	2006-Spring	2007-Spring	2008-Spring	2009-Spring	2010-Spring	2011-Spring	2012-Spring
Day 1 and 2	79000	22000	13600	7850	2470	5770	3080	23000	3100	5930	1260	1800
Day 3, 4, and 5	4700	7020	6580	38000	33000	57000	3060	5550	51900	8840	6060	20500

PCB results prior to 2000 were all less than detection limits. Starting in 2000 detection limits were lowered so that usable concentrations were available.

Note: For reference purposes, the ATSDR indicates that typical mean PCB concentrations in background soil are less than 100,000 pg/g

PAH	NYSDEC SCO ¹ unrestricted	NYSDEC SCO ² restricted	EPA screening level ³	NYS Rural soil survey ⁴	Tox. ⁵ Profile Rural soil background	Tox. ⁶ Profile Agr. Soil background	Tox. ⁷ Profile Urban soil background
Napthalene	12,000	100,000	3,900	17-24	NA	NA	NA
Acenaphthylene	100,000	100,000	3,400,000	110-500	NA	5	NA
Acenaphthene	20,000	100,000	NA	150	1.7	6	NA
Fluorene	30,000	100,000	2,300,000	580	NA	9.7	NA
Phenanthrene	100,000	100,000	NA	8,500	30	48-140	NA
Anthracene	100,000	100,000	17,000,000	620	NA	11-13	NA
Fluoranthene	100,000	100,000	2,300,000	7,400	0.3-40	120-210	200-166,000
Pyrene	100,000	100,00	1,700,000	8,700	1-19.7	99-150	145-147,000
Benzo(A)Anthracene	1,000	1,000	150	2,900	5-20	56-110	169-59,000
Chrysene	1,000	1,000	15,000	1,300	38.3	78-120	251-640
Benzo(B,K)Fluoranthene	1,000	1,000	150-1500	1,500-3,300	10-110	58-250	15,000-62,000
Benzo(E)Pyrene	NA	NA	NA	NA	NA	53-130	60-14,000
Benzo(A)Pyrene	1,000	1,000	15	2,400	2-1,300	4.6-900	165-220
Perylene	NA	NA	NA	8,700	NA	NA	NA
Dibenzo(A,H)Anthracene	330	330	15	NA	NA	NA	NA
Indeno(1,2,3-CD)Pyrene	500	500	150	660	10-15	63-100	8,000-61,000
Benzo(G,H,I)Perylene	100,000	100,000	NA	630	10-70	66	900-47,000
2-Methylnaphthalene	NA	NA	310,000	NA	NA	NA	NA
2-Chloronaphthalene	NA	NA	NA	NA	NA	NA	NA

Sources:

1,2. New York State Department of Environmental Soil Cleanup Objectives, 9/06. Unrestricted use accounts for the use of the land for raising livestock.

3. USEPA residential soil screening levels (SSL's), September, 2008/

4. NYS Rural Soil Survey, NYSDEC, 2005.

5,6,7. Agency for Toxic Substances and Disease Registry (ATSDR), Toxicological Profiles, 1995/